



RECEIVED

OCT 31 2003

TECH CENTER 1600/2900

Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.


.....

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date:

Signature:

Boston, 29 September 2003



Thomas Vorup Jensen, Ph.D.

OCT 31 2003

TECH CENTER 1600/2900



Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odcense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaebler heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date:

Signature:

2/4/97

Wilhelm Schwaebler

Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group, including research students Søren Hansen and Steen B. Laursen, showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....

To the best of my knowledge the above is a true account of the identification of MASP-2.


Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date: 22/9-03

Signature:

Steen B. Laurson

STEEN B. LAURSEN



Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date: 3 April 1997

Signature: *Kenneth Reid*



Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date: 14th April 1997

Signature:


A.C. WILLIS



Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting belong to Steffen Thiel and Jens Chr. Jensenius

Date: 21/11/96 .

Signature:

Discovery of MASP-2



The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group, including research students Søren Hansen and Steen B. Laursen, showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date: 14/9 - 2003
Odense, DK

Signature:

Discovery of MASP-2



The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....


To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting belong to Steffen Thiel and Jens Chr. Jensenius

Date:

29/11 1996

Signature:

A handwritten signature in cursive script, appearing to read "Uffe Holmskov".



Discovery of MASP-2

The protein now designated MASP-2 was isolated and identified in Aarhus by Steffen Thiel and Jens Chr. Jensenius through its reaction with a chicken antibody.

The N-terminal amino acids of the protein was sequenced in Oxford by Anthony C. Willis from blots on PVDF provided by the Aarhus group. Antibody against a synthetic peptide representing the N-terminal 20 amino acids reacted with the protein as well as with a polypeptide of 20 kDa. The 20 kDa peptide was N-terminally sequenced by ACW and found to be identical to the N-terminal sequence of the 50 kDa protein. Further, ACW sequenced some internal peptides of the protein and the 20 kDa form. These sequences established the protein as being related to MASP-1, C1r and C1s. The sequences also established the 20 kDa peptide as being a truncated form of the 50 kDa protein.

The Aarhus group showed the protein as well as the truncated form to be associated with mannan-binding lectin. Uffe Holmskov, Odense, was involved in this finding through ongoing discussions.

In Aarhus graduate student Thomas Vorup Jensen generated by RT-PCR a 300 bp cDNA fragment from liver RNA using oligonucleotides deduced from the peptide sequences. The amino acid sequence deduced from the 300 bp oligonucleotide encompassed the peptides used for constructing the oligonucleotides as well as another of the sequenced peptides. The derived sequence further emphasized the similarity of the new protein to MASP (MASP-1) and also the relationship to C1r and C1s.

TVJ working in Kenneth B. M. Reid's laboratory in Oxford with guidance of KBMR and Paul Eggleton used the 300 bp as probe for cloning cDNA from liver libraries and isolated a 1.8 kb clone. The sequence revealed the presence of the serine protease domain, although the C-terminal part of this was missing. Also missing (compared to the later obtained full length clone) was the 2nd C1r/s domain.

Wilhelm Schwaeble heard of the project and offered his help with isolating and sequencing further clones. The aim of this was to establish the full sequence of the mRNA and thereby also the deduced amino acid sequence. This work would also allow for the synthesis by recombinant technique of the recombinant protein. The synthesis of the recombinant protein as well as of MASP-1 and possibly also MBL was allocated to TVJ to be part of his PhD project aimed at studying the structure and function of the MBL/MASP complex. This work was designed to be carried out in collaboration with KBMR and Robert B. Sim in Oxford as well as at the Aarhus laboratory.

With the use of the 300 bp RT-PCR generated probe, the cloning and sequencing was successfully carried out in WS's laboratory in Leicester, largely by graduate student Cordula Stover. Three different cDNA clones were sequenced. One (A) represents the full length was found to encode a protein of 622 amino acids showing about 40% sequence identity to C1r, C1s and MASP-1. Another clone (C) represented the N-terminal part of clone A with additional four amino acids not found in A. The size of C (ORF of 540 bp) agrees with it representing mRNA encoding the truncated form of MASP-2. The third clone (B) has a 5' end almost identical to C but in addition an ORF of 558 bp with no similarity to A.

.....

To the best of my knowledge the above is a true account of the identification of MASP-2. Being coauthor of the first publication on MASP-2 I further hereby state that I concede that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong to Steffen Thiel and Jens Chr. Jensenius

Date: 14th April 1997 Signature: Kenneth BM Reid.

Kim Palle Jensen



EX. A

From: Jens Christian Jensenius [j.c.jensenius@immunology.au.dk]
Sent: 7. oktober 2003 10:20
To: Cordula Stover
Subject: Re: signature

Dear Cordula,

Thank you for your mail. I certainly think that you deserve, and wish for you, a successful career. While I shall not lay claim to being an expert on these matters, I have been involved on a number of occasions, and my experience is that IPR is rather different from allocating scientific credit, where emphasis, contrary to IPR, is to a large extent based on the invested hard work. It is entirely a different world.

The best thing for me now is probably to leave the matter at our end to the patent agency.

Best wishes,

Jens

Dear Jens.

Yes, I have problems signing the paper on MASP-2 discovery thereby conceding that the rights of commercial exploitation and patenting resulting from the first publication (Nature April 1997) belong solely to you and Steffen.

I am on a successful career and have the obligation to myself to make use of any source I have co-explored.

With kindest regards

Cordula

on 3/10/03 16:51, Jens Christian Jensenius at j.c.jensenius@immunology.au.dk wrote:

Dear Cordula,

Sorry to bother you again. Please, tell me if you have problems signing the paper on MASP-2 discovery.

If, for some reason you decide not to sign, I shall have to sign a statement on your situation - being a PhD student with Wilhelm - etc etc - and this will be legally valid - can obviously be challenged - BUT the agency and the lawyers seem to prefer all documents being the same. Your signature is the only one missing.

Thus it will be convenient if you could, please, sign and return it to the the agency ASAP.

Best wishes, Jens

07-10-03

Fax: +45 8619 6128

privat

+45 8612 7124

+45 6611 5003

+45 9848 8947